

IBUPROFEN- ibuprofen tablet
WALGREENS

612R Walgreens 0363-9612 Ibuprofen 200mg Caplets

DRUG FACTS

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

temporarily relieves minor aches and pain due to :

- backache
- headache
- menstrual cramps
- minor pain of arthritis
- muscular aches
- the common cold
- toothache

temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergy reaction, especially in people allergic to aspirin.

Symptoms may include:

- asthma (wheezing)
- blisters
- facial swelling
- hives
- rash
- shock
- skin reddening

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have bad stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

you have symptoms of heart problem or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of the body
- slurred speech
- leg swelling

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

redness or swelling is present in the painful area

any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

Directions

do not take more than directed

the smallest effective dose should be used

adults and children 12 years and over:

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor

children under 12 years:

- ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

Questions or comments? call **1-877-290-4008**

50 COUNT 612R

100 COUNT 612R

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., Medical Consumer Healthcare Division, owner of the registered trademark Motrin®.

Lot No.:
Exp Date:

DISTRIBUTED BY: WALGREEN CO.

Active ingredient (in each capsule)	Purpose
100 mg of ampicillin sodium	Antibiotic
250 mg of ampicillin sodium	Antibiotic
500 mg of ampicillin sodium	Antibiotic
750 mg of ampicillin sodium	Antibiotic
1000 mg of ampicillin sodium	Antibiotic
1250 mg of ampicillin sodium	Antibiotic
1500 mg of ampicillin sodium	Antibiotic
1750 mg of ampicillin sodium	Antibiotic
2000 mg of ampicillin sodium	Antibiotic
2250 mg of ampicillin sodium	Antibiotic
2500 mg of ampicillin sodium	Antibiotic
2750 mg of ampicillin sodium	Antibiotic
3000 mg of ampicillin sodium	Antibiotic
3250 mg of ampicillin sodium	Antibiotic
3500 mg of ampicillin sodium	Antibiotic
3750 mg of ampicillin sodium	Antibiotic
4000 mg of ampicillin sodium	Antibiotic
4250 mg of ampicillin sodium	Antibiotic
4500 mg of ampicillin sodium	Antibiotic
4750 mg of ampicillin sodium	Antibiotic
5000 mg of ampicillin sodium	Antibiotic
5250 mg of ampicillin sodium	Antibiotic
5500 mg of ampicillin sodium	Antibiotic
5750 mg of ampicillin sodium	Antibiotic
6000 mg of ampicillin sodium	Antibiotic
6250 mg of ampicillin sodium	Antibiotic
6500 mg of ampicillin sodium	Antibiotic
6750 mg of ampicillin sodium	Antibiotic
7000 mg of ampicillin sodium	Antibiotic
7250 mg of ampicillin sodium	Antibiotic
7500 mg of ampicillin sodium	Antibiotic
7750 mg of ampicillin sodium	Antibiotic
8000 mg of ampicillin sodium	Antibiotic
8250 mg of ampicillin sodium	Antibiotic
8500 mg of ampicillin sodium	Antibiotic
8750 mg of ampicillin sodium	Antibiotic
9000 mg of ampicillin sodium	Antibiotic
9250 mg of ampicillin sodium	Antibiotic
9500 mg of ampicillin sodium	Antibiotic
9750 mg of ampicillin sodium	Antibiotic
10000 mg of ampicillin sodium	Antibiotic

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

- temporarily relieves minor aches and pains due to:
 - backache ● headache ● menstrual cramps
 - minor pain of arthritis ● muscular aches
 - the common cold ● toothache
- temporarily reduces fever

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

Symptoms may include:

- asthma (wheezing)
- hives
- rash
- shock
- skin peeling
- itchy, watery, red, swollen eyes
- itchy, watery, red, swollen mouth
- itchy, watery, red, swollen throat
- itchy, watery, red, swollen nose
- itchy, watery, red, swollen lips
- itchy, watery, red, swollen tongue
- itchy, watery, red, swollen skin
- itchy, watery, red, swollen hair
- itchy, watery, red, swollen nails
- itchy, watery, red, swollen teeth
- itchy, watery, red, swollen gums
- itchy, watery, red, swollen jaw
- itchy, watery, red, swollen chin
- itchy, watery, red, swollen neck
- itchy, watery, red, swollen shoulders
- itchy, watery, red, swollen arms
- itchy, watery, red, swollen hands
- itchy, watery, red, swollen fingers
- itchy, watery, red, swollen toes
- itchy, watery, red, swollen feet
- itchy, watery, red, swollen ankles
- itchy, watery, red, swollen legs
- itchy, watery, red, swollen hips
- itchy, watery, red, swollen buttocks
- itchy, watery, red, swollen groin
- itchy, watery, red, swollen perineum
- itchy, watery, red, swollen anus
- itchy, watery, red, swollen vagina
- itchy, watery, red, swollen penis
- itchy, watery, red, swollen testicles
- itchy, watery, red, swollen scrotum
- itchy, watery, red, swollen urethra
- itchy, watery, red, swollen bladder
- itchy, watery, red, swollen stomach
- itchy, watery, red, swollen intestines
- itchy, watery, red, swollen liver
- itchy, watery, red, swollen gallbladder
- itchy, watery, red, swollen pancreas
- itchy, watery, red, swollen spleen
- itchy, watery, red, swollen lungs
- itchy, watery, red, swollen heart
- itchy, watery, red, swollen kidneys
- itchy, watery, red, swollen bladder
- itchy, watery, red, swollen uterus
- itchy, watery, red, swollen ovaries
- itchy, watery, red, swollen fallopian tubes
- itchy, watery, red, swollen cervix
- itchy, watery, red, swollen vagina
- itchy, watery, red, swollen penis
- itchy, watery, red, swollen testicles
- itchy, watery, red, swollen scrotum
- itchy, watery, red, swollen urethra
- itchy, watery, red, swollen bladder
- itchy, watery, red, swollen stomach
- itchy, watery, red, swollen intestines
- itchy, watery, red, swollen liver
- itchy, watery, red, swollen gallbladder
- itchy, watery, red, swollen pancreas
- itchy, watery, red, swollen spleen
- itchy, watery, red, swollen lungs
- itchy, watery, red, swollen heart
- itchy, watery, red, swollen kidneys

Normal healing warning: The product contains an NSAID, which may cause some stomach bleeding. The chance of this is higher if you: average 60 or older have had stomach issues in the past are taking blood-thinning medications, or a steroid drug have had other drug interactions, prescription or over-the-counter NSAIDs (aspirin, ibuprofen, naproxen, or others) take 3 or more alcoholic drinks every day while taking the product.

Take more or for a longer time than directed.

- you are taking a diuretic
- Ask a doctor or pharmacist before use if you are
 - taking aspirin for heart attack or stroke, because aspirin may decrease the level of aspirin
 - under a doctor's care for any serious condition
 - taking any other drug

When using this product

- Use with food or milk. Stomach upset occurs
- Don't use an alkali tablet if:
 - you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have blood or black stools
 - have stomach pain that does not get better
 - you have symptoms of heart and kidney trouble
 - feel pain
 - trouble breathing

Heart attack and stroke warning: MSADs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning signs to you
- you have a history of stomach problems, such as heartburn

- weakness in one part or side of the body
- slurred speech ● big swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area

■ any new symptoms appear

<ul style="list-style-type: none"> ■ do not take more than five doses ■ the smallest effective dose should be used 	adults and children 12 years and over: <ul style="list-style-type: none"> ■ take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor
	children under 12 years: <ul style="list-style-type: none"> ■ ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Inactive ingredients (continued)
colloidal silicon dioxide, croscarmellose sodium, FD & C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

ORIG00000
REV-0818
612F

NDC 0363-9612-01

Ibuprofen 200

IBUPROFEN TABLETS USP, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

CAPLETS

100 CAPLETS**
(*CAPSULE-SHAPED TABLETS)

SEE NEW WARNINGS INFORMATION

ACTUAL COSTS


E827

Varnish/Print Omit Area

Varnish
Omit

225 COUNT 612R

<p>Drug Facts (continued)</p> <p>It is an aspirin. You have overdosed an aspirin because you took too many tablets. You have overdosed on ibuprofen or you either gave an aspirin or ibuprofen to your child or their health is in jeopardy.</p> <p>Other important information:</p> <ul style="list-style-type: none"> ● Aspirin: A fatal illness called Reye's syndrome can develop in children who take aspirin. This illness can be fatal. Do not give aspirin to children. ● Ibuprofen: Ibuprofen may irritate the stomach and cause bleeding. Do not give ibuprofen to children. ● Aspirin and ibuprofen: Do not give your child both aspirin and ibuprofen together. This can cause bleeding in the stomach. ● Other medicines: Do not give your child any other medicines without talking to a doctor first. Some medicines can interact with aspirin or ibuprofen. 	<p>What you should know:</p> <ul style="list-style-type: none"> ● Aspirin: Do not give your child aspirin. It can cause a fatal illness called Reye's syndrome. This illness can be fatal. ● Ibuprofen: Do not give your child ibuprofen. It can cause bleeding in the stomach. ● Aspirin and ibuprofen: Do not give your child both aspirin and ibuprofen together. This can cause bleeding in the stomach. ● Other medicines: Do not give your child any other medicines without talking to a doctor first. Some medicines can interact with aspirin or ibuprofen. 	<p>What you should know:</p> <ul style="list-style-type: none"> ● Aspirin: Do not give your child aspirin. It can cause a fatal illness called Reye's syndrome. This illness can be fatal. ● Ibuprofen: Do not give your child ibuprofen. It can cause bleeding in the stomach. ● Aspirin and ibuprofen: Do not give your child both aspirin and ibuprofen together. This can cause bleeding in the stomach. ● Other medicines: Do not give your child any other medicines without talking to a doctor first. Some medicines can interact with aspirin or ibuprofen. 	<p>What you should know:</p> <ul style="list-style-type: none"> ● Aspirin: Do not give your child aspirin. It can cause a fatal illness called Reye's syndrome. This illness can be fatal. ● Ibuprofen: Do not give your child ibuprofen. It can cause bleeding in the stomach. ● Aspirin and ibuprofen: Do not give your child both aspirin and ibuprofen together. This can cause bleeding in the stomach. ● Other medicines: Do not give your child any other medicines without talking to a doctor first. Some medicines can interact with aspirin or ibuprofen. 	<p>What you should know:</p> <ul style="list-style-type: none"> ● Aspirin: Do not give your child aspirin. It can cause a fatal illness called Reye's syndrome. This illness can be fatal. ● Ibuprofen: Do not give your child ibuprofen. It can cause bleeding in the stomach. ● Aspirin and ibuprofen: Do not give your child both aspirin and ibuprofen together. This can cause bleeding in the stomach. ● Other medicines: Do not give your child any other medicines without talking to a doctor first. Some medicines can interact with aspirin or ibuprofen. 	<p>What you should know:</p> <ul style="list-style-type: none"> ● Aspirin: Do not give your child aspirin. It can cause a fatal illness called Reye's syndrome. This illness can be fatal. ● Ibuprofen: Do not give your child ibuprofen. It can cause bleeding in the stomach. ● Aspirin and ibuprofen: Do not give your child both aspirin and ibuprofen together. This can cause bleeding in the stomach. ● Other medicines: Do not give your child any other medicines without talking to a doctor first. Some medicines can interact with aspirin or ibuprofen. 	<p>What you should know:</p> <ul style="list-style-type: none"> ● Aspirin: Do not give your child aspirin. It can cause a fatal illness called Reye's syndrome. This illness can be fatal. ● Ibuprofen: Do not give your child ibuprofen. It can cause bleeding in the stomach. ● Aspirin and ibuprofen: Do not give your child both aspirin and ibuprofen together. This can cause bleeding in the stomach. ● Other medicines: Do not give your child any other medicines without talking to a doctor first. Some medicines can interact with aspirin or ibuprofen. 	<p>What you should know:</p> <ul style="list-style-type: none"> ● Aspirin: Do not give your child aspirin. It can cause a fatal illness called Reye's syndrome. This illness can be fatal. ● Ibuprofen: Do not give your child ibuprofen. It can cause bleeding in the stomach. ● Aspirin and ibuprofen: Do not give your child both aspirin and ibuprofen together. This can cause bleeding in the stomach. ● Other medicines: Do not give your child any other medicines without talking to a doctor first. Some medicines can interact with aspirin or ibuprofen. 	<p>What you should know:</p> <ul style="list-style-type: none"> ● Aspirin: Do not give your child aspirin. It can cause a fatal illness called Reye's syndrome. This illness can be fatal. ● Ibuprofen: Do not give your child ibuprofen. It can cause bleeding in the stomach. ● Aspirin and ibuprofen: Do not give your child both aspirin and ibuprofen together. This can cause bleeding in the stomach. ● Other medicines: Do not give your child any other medicines without talking to a doctor first. Some medicines can interact with aspirin or ibuprofen.
---	--	--	--	--	--	--	--	--




Compare to Motrin® IB
Caplets active ingredient!

NDC 0393-961 2-26

Ibuprofen 200

IBUPROFEN TABLETS USP, 200 mg /
PAIN RELIEVER / FEVER REDUCER (NSAID)

CAPLETS



225 CAPLETS**
(**CAPSULE-SHAPED TABLETS)

ACTUAL SIZE

SEE NEW WARNINGS INFORMATION

THIS PRODUCT FOR PAIN RELIEF WITHOUT TYPICAL CHILDREN

COMPARE TO MOTRIN® IB THAT YOUR PARENTS
GAVE YOU TO TAKE CARE OF THE MOUTH OF
THE BOTTLE LICK, TOOK AROUND OR DRINKS

Compare to Motrin® IB
Ibuprofen 200 mg Tablets

Drugs Facts
Active ingredient (in each caplet):
Ibuprofen 200 mg (NSAID)
*most brands with information drug

Uses
Temporarily relieve minor aches and pains due to:
• headache • teething • menstrual cramps • minor pain of
arthritis • muscular aches • the common cold • tooth ache
• temporarily reduce fever

Warnings
Allergy Alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
• difficulty breathing • hives • itchy skin • rash • swelling
• asthma (wheezing) • dizziness • facial swelling
• nausea • vomiting • skin rash • skin redness • skin allergic
reaction occurs, stop use and seek medical help right away.
Stomach Bleeding Warning: This product contains an NSAID,
which may cause some stomach bleeding. The chance is higher if
you take aspirin, 80 mg or more, at the same time.
• Have had stomach ulcers or bleeding problems
• Take a blood-thinning (anticoagulant) or steroid drug
• Take other drugs (including prescription or nonprescription
NSAIDs, aspirin, ibuprofen, naproxen, or others)
• Have 3 or more alcoholic drinks every day while taking this
product • Take more for a longer time than directed
Heart Attack and Stroke Warning: NSAIDs, except aspirin, increase
the risk of heart attack, heart failure, and stroke. These can be fatal.
The risk is higher if you use more than directed or for longer than
directed.

Drug Facts (continued from label)
The product is not a narcotic and is not habit forming. Johnson & Johnson Consumer Inc.,
Mallinckrodt Consumer Health Division, One of the Registered Trademarks of Mallinckrodt

US: TROST OF THE WALGREEN CO.
100% SATISFACTION GUARANTEED
NATIONWIDE 800-368-6789

LOT No.: 311917-204819
Exp. Date:

MADE IN INDIA
ITEM# 2203-85
400000-0000-X

8/28 REV 1-18
OR03000

[illegible]

IBUPROFEN

ibuprofen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-9612
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	orange	Score	no score
Shape	CAPSULE	Size	10mm
Flavor		Imprint Code	120
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9612-05	50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/11/2018	
2	NDC:0363-9612-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/11/2018	
3	NDC:0363-9612-26	225 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/11/2018	
4	NDC:0363-9612-50	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/11/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091237	12/11/2018	

Labeler - WALGREENS (008965063)

Registrant - TIME CAP LABORATORIES, INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(0363-9612)

Revised: 11/2021

WALGREENS